

Multiscale simulation of polymeric fluids using massively parallel computers

Finite Elements in Flow Problems 2017 – Rome, Italy

April 5th – 7th

Alexander Rüttgers

German Aerospace Center (DLR)

Simulation and Software Technology

Cologne, Germany




Knowledge for Tomorrow



OUTLINE

- 1 MODELING EQUATIONS FOR MULTISCALE APPROACH
- 2 NUMERICAL DISCRETIZATION
- 3 SIMULATION RESULT: 3D CONTRACTION FLOW
- 4 COMPUTATIONAL COMPLEXITY

MODELING OF POLYMERIC FLUIDS




Two different modeling approaches:

overview ¹	macroscopic	multiscale
cost	low	high
modeling accuracy	low	high
drawbacks	num. instabilities	


¹ Le Bris and Lelièvre 2009

MICROSCALE MODELING



- Position vector \mathbf{x} in flow space $\mathcal{O} \subset \mathbb{R}^3$.
- $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ in configuration space $\mathcal{D} \subset \mathbb{R}^{3N}$.

MICROSCALE MODELING




- Position vector \mathbf{x} in flow space $\mathcal{O} \subset \mathbb{R}^3$.
- $\mathbf{q} = (q_1, \dots, q_N)$ in configuration space $\mathcal{D} \subset \mathbb{R}^{3N}$.

Description of a polymer ensemble

- **Stochastic approach:**¹ random field $\mathbf{Q} = (Q_1, \dots, Q_N)$ with $\mathbf{Q} : (\mathbf{x}, t) \in \mathcal{O} \times \mathcal{T} \mapsto \mathbf{Q}(\mathbf{x}, t)$ as \mathcal{D} -valued random variable.

¹Laso and Öttinger 1993

MICROSCALE MODELING



- Position vector \mathbf{x} in flow space $\mathcal{O} \subset \mathbb{R}^3$.
- $\mathbf{q} = (q_1, \dots, q_N)$ in configuration space $\mathcal{D} \subset \mathbb{R}^{3N}$.

Description of a polymer ensemble

- **Stochastic approach:**¹ random field $\mathbf{Q} = (\mathbf{Q}_1, \dots, \mathbf{Q}_N)$ with $\mathbf{Q} : (\mathbf{x}, t) \in \mathcal{O} \times \mathcal{T} \mapsto \mathbf{Q}(\mathbf{x}, t)$ as \mathcal{D} -valued random variable.
- **Fokker-Planck ansatz:**² probability density function of field \mathbf{Q}
 - $\psi : (\mathbf{x}, \mathbf{q}, t) \in \mathcal{O} \times \mathcal{D} \times \mathcal{T} \subset \mathbb{R}^{3N+4} \mapsto \psi(\mathbf{x}, \mathbf{q}, t) \in \mathbb{R}^+$.
 - $\int_{\mathcal{D}} \psi(\mathbf{x}, \mathbf{q}, t) d\mathbf{q} = 1$ for all $(\mathbf{x}, t) \in \mathcal{O} \times \mathcal{T}$.

¹Laso and Öttinger 1993 , ²Lozinski and Chauvière 2003

COUPLING OF MICRO- AND MACROSCOPIC SCALE

- Elastic fluid behavior modeled with macroscopic stress tensor $\tau_p : (\mathbf{x}, t) \in \mathcal{O} \times \mathcal{T} \mapsto \tau_p(\mathbf{x}, t) \in \mathbb{R}^{3 \times 3}$.

COUPLING OF MICRO- AND MACROSCOPIC SCALE

- Elastic fluid behavior modeled with macroscopic stress tensor $\boldsymbol{\tau}_p : (\mathbf{x}, t) \in \mathcal{O} \times \mathcal{T} \mapsto \boldsymbol{\tau}_p(\mathbf{x}, t) \in \mathbb{R}^{3 \times 3}$.
- **Kramers'**¹ relation for stress tensor

$$\boldsymbol{\tau}_p(\mathbf{x}, t) = \begin{pmatrix} \tau_{xx} & \tau_{xy} & \tau_{xz} \\ \tau_{yx} & \tau_{yy} & \tau_{yz} \\ \tau_{zx} & \tau_{zy} & \tau_{zz} \end{pmatrix}(\mathbf{x}, t) = C \sum_{i=1}^N (\mathbb{E}[\mathbf{Q}_i(\mathbf{x}, t) \otimes \mathbf{F}(\mathbf{Q}_i(\mathbf{x}, t))] - \mathbf{Id})$$

- expectation $\mathbb{E}[\cdot] = \int_{\mathcal{D}} \cdot \psi(\mathbf{x}, \mathbf{q}, t) d\mathbf{q}$ in configuration space \mathcal{D} .
- spring force $\mathbf{F} : \mathcal{D}_i \subset \mathbb{R}^3 \rightarrow \mathbb{R}^3$.

¹Kramers 1944

COUPLING OF MICRO- AND MACROSCOPIC SCALE

- Elastic fluid behavior modeled with macroscopic stress tensor
 $\tau_{\mathbf{p}} : (\mathbf{x}, t) \in \mathcal{O} \times \mathcal{T} \mapsto \tau_{\mathbf{p}}(\mathbf{x}, t) \in \mathbb{R}^{3 \times 3}$.
- **Kramers'**¹ relation for stress tensor

$$\tau_{\mathbf{p}}(\mathbf{x}, t) = \begin{pmatrix} \tau_{xx} & \tau_{xy} & \tau_{xz} \\ \tau_{yx} & \tau_{yy} & \tau_{yz} \\ \tau_{zx} & \tau_{zy} & \tau_{zz} \end{pmatrix}(\mathbf{x}, t) = C \sum_{i=1}^N (\mathbb{E}[\mathbf{Q}_i(\mathbf{x}, t) \otimes \mathbf{F}(\mathbf{Q}_i(\mathbf{x}, t))] - \mathbf{Id})$$

- expectation $\mathbb{E}[\cdot] = \int_{\mathcal{D}} \cdot \psi(\mathbf{x}, \mathbf{q}, t) d\mathbf{q}$ in configuration space \mathcal{D} .
- spring force $\mathbf{F} : \mathcal{D}_i \subset \mathbb{R}^3 \rightarrow \mathbb{R}^3$.
- Various spring force models \mathbf{F} in the literature
 - **Hooke** model : $\mathbf{F}(\mathbf{Q}_i) = \mathbf{Q}_i$ (linear)
 - **FENE** model² : $\mathbf{F}(\mathbf{Q}_i) = \frac{\mathbf{Q}_i}{1 - \|\mathbf{Q}_i\|^2/b}$ with $\|\mathbf{Q}_i\|^2 \leq b \in \mathbb{R}^+$ (nonlinear)
 - **CPAIL** model³: $\mathbf{F}(\mathbf{Q}_i) = \frac{1 - \|\mathbf{Q}_i\|^2/(3b)}{1 - \|\mathbf{Q}_i\|^2/b} \mathbf{Q}_i$ with $\|\mathbf{Q}_i\|^2 \leq b \in \mathbb{R}^+$ (nonlinear)

¹Kramers 1944, ²Warner 1972, ³Cohen 1991

STOCHASTIC MULTISCALE SYSTEM

$$\frac{D\mathbf{U}(\mathbf{x}, t)}{Dt} = -\nabla P(\mathbf{x}, t) + \frac{\beta}{Re} \Delta \mathbf{U}(\mathbf{x}, t) + \frac{1}{Re} \nabla \cdot \boldsymbol{\tau}_p(\mathbf{x}, t) \quad (1)$$

$$\nabla \cdot \mathbf{U}(\mathbf{x}, t) = 0 \quad (2)$$

$$d\mathbf{Q}(\mathbf{x}, t) = \left[-\mathbf{U}(\mathbf{x}, t) \nabla \mathbf{Q}(\mathbf{x}, t) + (\nabla \mathbf{U}(\mathbf{x}, t))^T \mathbf{Q}(\mathbf{x}, t) - \frac{1}{4De} \mathbf{A} \cdot \mathbf{F}(\mathbf{Q}(\mathbf{x}, t)) \right] dt + \sigma d\mathbf{W}(t) \quad (3)$$

$$\boldsymbol{\tau}_p(\mathbf{x}, t) = C \sum_{i=1}^N (\mathbb{E}[\mathbf{Q}_i(\mathbf{x}, t) \otimes \mathbf{F}(\mathbf{Q}_i(\mathbf{x}, t))] - \mathbf{Id}) . \quad (4)$$

for the unknown random fields \mathbf{U} , P , \mathbf{Q} , $\boldsymbol{\tau}_p$,
dimensionless parameters $De, Re, \beta, \sigma \in \mathbb{R}^+$
+ initial and boundary conditions.

(1)+(2) Navier-Stokes equations (macroscopic)

(3) Stochastic PDE (microscopic)

(4) upscaling from micro- to macroscopic scale

STOCHASTIC APPROACH FOR MICROSCALE^{1,2}

- Approximation of random field $\mathbf{Q}(\mathbf{x}_k, t)$ at discrete points \mathbf{x}_k with $3N$ -dimensional **samples** $\mathbf{Q}^{(j)}(\mathbf{x}_k, t) \sim \psi(\mathbf{x}_k, \cdot, t)$ for $j = 1, \dots, M_s$.

¹Hulsen et al. 1997, ²Laso and Öttinger 1993

STOCHASTIC APPROACH FOR MICROSCALE^{1,2}

- Approximation of random field $\mathbf{Q}(\mathbf{x}_k, t)$ at discrete points \mathbf{x}_k with $3N$ -dimensional **samples** $\mathbf{Q}^{(j)}(\mathbf{x}_k, t) \sim \psi(\mathbf{x}_k, \cdot, t)$ for $j = 1, \dots, M_s$.
- Monte Carlo approximation of expectation

$$\begin{aligned}\tau_{\mathbf{p}}(\mathbf{x}_k, t) &= C \sum_{i=1}^N \left(\mathbb{E}[\mathbf{Q}_i(\mathbf{x}_k, t) \otimes \mathbf{F}_i(\mathbf{Q}_i(\mathbf{x}_k, t))] - \mathbf{Id} \right) \\ &\approx C \sum_{i=1}^N \left(\frac{1}{M_s} \sum_{j=1}^{M_s} \mathbf{Q}_i^{(j)}(\mathbf{x}_k, t) \otimes \mathbf{F}_i(\mathbf{Q}_i^{(j)}(\mathbf{x}_k, t)) - \mathbf{Id} \right).\end{aligned}$$

¹Hulsen et al. 1997, ²Laso and Öttinger 1993

STOCHASTIC APPROACH FOR MICROSCALE^{1,2}

- Approximation of random field $\mathbf{Q}(\mathbf{x}_k, t)$ at discrete points \mathbf{x}_k with $3N$ -dimensional **samples** $\mathbf{Q}^{(j)}(\mathbf{x}_k, t) \sim \psi(\mathbf{x}_k, \cdot, t)$ for $j = 1, \dots, M_s$.
- Monte Carlo approximation of expectation

$$\begin{aligned}\tau_{\mathbf{p}}(\mathbf{x}_k, t) &= C \sum_{i=1}^N \left(\mathbb{E}[\mathbf{Q}_i(\mathbf{x}_k, t) \otimes \mathbf{F}_i(\mathbf{Q}_i(\mathbf{x}_k, t))] - \mathbf{Id} \right) \\ &\approx C \sum_{i=1}^N \left(\frac{1}{M_s} \sum_{j=1}^{M_s} \mathbf{Q}_i^{(j)}(\mathbf{x}_k, t) \otimes \mathbf{F}_i(\mathbf{Q}_i^{(j)}(\mathbf{x}_k, t)) - \mathbf{Id} \right).\end{aligned}$$

- **Variance error** in $\tau_{\mathbf{p}}$ is of order $\mathcal{O}(M_s^{-1/2})$.


¹Hulsen et al. 1997, ²Laso and Öttinger 1993

STOCHASTIC APPROACH FOR MICROSCALE^{1,2}

- Approximation of random field $\mathbf{Q}(\mathbf{x}_k, t)$ at discrete points \mathbf{x}_k with $3N$ -dimensional **samples** $\mathbf{Q}^{(j)}(\mathbf{x}_k, t) \sim \psi(\mathbf{x}_k, \cdot, t)$ for $j = 1, \dots, M_s$.
- Monte Carlo approximation of expectation

$$\begin{aligned}\tau_p(\mathbf{x}_k, t) &= C \sum_{i=1}^N \left(\mathbb{E}[\mathbf{Q}_i(\mathbf{x}_k, t) \otimes \mathbf{F}_i(\mathbf{Q}_i(\mathbf{x}_k, t))] - \mathbf{Id} \right) \\ &\approx C \sum_{i=1}^N \left(\frac{1}{M_s} \sum_{j=1}^{M_s} \mathbf{Q}_i^{(j)}(\mathbf{x}_k, t) \otimes \mathbf{F}_i(\mathbf{Q}_i^{(j)}(\mathbf{x}_k, t)) - \mathbf{Id} \right).\end{aligned}$$

- **Variance error** in τ_p is of order $\mathcal{O}(M_s^{-1/2})$.




¹Hulsen et al. 1997, ²Laso and Öttinger 1993

TEMPORAL EVOLUTION OF MICROSCOPIC SCALE

- Density ψ only known at $t = 0$ for complex spring models.
- At $t = 0$ create stochastic samples using **rejection sampling**¹.


temporal evolution
of stoch. samples



¹von Neumann 1951



TEMPORAL EVOLUTION OF MICROSCOPIC SCALE

- Density ψ only known at $t = 0$ for complex spring models.
- At $t = 0$ create stochastic samples using **rejection sampling**¹.
- For $t > 0$: Semi-implicit **Euler-Maruyama** scheme with 1. order accuracy in time.



¹ von Neumann 1951



SIMULATION: 3D CONTRACTION FLOW



- Applications: injection moulding, polymer processing, ...
- Discretization in space with 3D flow solver *NaSt3dGPF*^{1,2,3} on a staggered grid of size M_g with center \mathbf{x}_k .

¹Griebel et al. 1998, ²Croce et al. 2009 ³ For non-profit use: <http://wissrech.ins.uni-bonn.de>

SIMULATION: 3D CONTRACTION FLOW



- Applications: injection moulding, polymer processing, ...
- Discretization in space with 3D flow solver *NaSt3dGPF*^{1,2,3} on a staggered grid of size M_g with center \mathbf{x}_k .
- Dimensionless characteristic units:

Reynolds number: $Re = \frac{2L\rho U}{\eta(\dot{\gamma})}$,

Deborah number: $De = \frac{\lambda U}{L}$

U average velocity, λ relaxation time,
 $\eta(\dot{\gamma})$ viscosity, $\dot{\gamma}$ shear rate, L characteristic length

¹Griebel et al. 1998, ²Croce et al. 2009 ³ For non-profit use: <http://wissrech.ins.uni-bonn.de>

SHEAR-THINNING FLUID IN 4 : 1 CONTRACTION

- Experimental measurements:¹
 - fluid: Glycerol + water + PAA (600ppm)
 - relaxation time: $\lambda = 32\text{s}$
 - experimental Deborah numbers: $De = 1, \dots, 200$




FIGURE: Visualization of contraction flow¹ with $Re = 2.37$, $De = 174$

¹Sousa, Coelho, Oliveira and Alves 2011

SHEAR-THINNING FLUID IN 4 : 1 CONTRACTION

- Experimental measurements:¹
 - fluid: Glycerol + water + PAA (600ppm)
 - relaxation time: $\lambda = 32\text{s}$
 - experimental Deborah numbers: $De = 1, \dots, 200$
- For high Deborah number flows:
 - large corner vortices occur
 - streamline divergence
 - inverted streamline rotation
- Macroscopic approaches often suffer from numerical instabilities^{2,3}.




FIGURE: Visualization of contraction flow¹ with $Re = 2.37$, $De = 174$

¹Sousa, Coelho, Oliveira and Alves 2011, ²Keunings 1986, ³Mangoubi et al. 2009

3D SIMULATION RESULTS¹

- 8 multiscale simulations.

Simulation parameters	
Deborah number De	24.1, 108, 157
spring model	FENE chain
spring segments N	1, 3, 5
resolution M_g	$380 \times 64 \times 64$
samples per cell M_s	1200

¹partial results in Griebel and R. 2014

3D SIMULATION RESULTS¹

- 8 multiscale simulations.
- All elastic effects reproduced in simulation.
- No stability issues occurred.

Simulation parameters		
Deborah number De	24.1, 108, 157	
spring model	FENE chain	
spring segments N	1, 3, 5	
resolution M_g	$380 \times 64 \times 64$	
samples per cell M_s	1200	




FIGURE: Simulation of 5-segment chain with $De = 157$.

¹partial results in Griebel and R. 2014

3D SIMULATION RESULTS¹

- 8 multiscale simulations.
- All elastic effects reproduced in simulation.
- No stability issues occurred.

Simulation parameters

Deborah number De	24.1, 108, 157
spring model	FENE chain
spring segments N	1, 3, 5
resolution M_g	$380 \times 64 \times 64$
samples per cell M_s	1200





FIGURE: Simulation of 5-segment chain with $De = 157$.


¹partial results in Griebel and R. 2014

COMPARISON OF EXPERIMENT AND SIMULATION


- **Velocity profiles** compared on the channel's centerline (a).
- **Inverted 3D streamline rotation** compared to Newtonian flow (b+c).
- Simulation results correspond with experimental measurements.



(a) velocity profile $De=24$



(b) streamlines $De = 1.0$



(c) streamlines $De=157$

COMPLEXITY OF MULTISCALE SIMULATIONS

Clusters for multiscale simulations

	Atacama	JUROPA
# cores	1,248 CPUs	17,664 CPUs
memory	4,992 GB	52,992 GB
Linpack	≈ 21 TFlops/s	207 TFlops/s
installation	Mar 2014	access: 11/2013 - 10/2014

Institute for Numerical Simulation, University of Bonn

Jülich Supercomputing Centre, Jülich Research Centre



(a) INS cluster



(b) Juropa racks

COMPLEXITY OF MULTISCALE SIMULATIONS

- M_g grid cells in space,
- M_s samples per grid cell,
- spring model with N segments.

COMPLEXITY OF MULTISCALE SIMULATIONS

- M_g grid cells in space,
- M_s samples per grid cell,
- spring model with N segments.

Complexity of stochastic microscale	
spatial grid M_g	$380 \times 64 \times 64$
samples per cell M_s	1200
total samples	$1.87 \cdot 10^9$
spring segments N	5
sample dimensionality	15-dim
storage	208 GB
computing time (256 CPUs)	7-8 weeks

COMPLEXITY OF MULTISCALE SIMULATIONS

- M_g grid cells in space,
- M_s samples per grid cell,
- spring model with N segments.

Complexity of stochastic microscale	
spatial grid M_g	$380 \times 64 \times 64$
samples per cell M_s	1200
total samples	$1.87 \cdot 10^9$
spring segments N	5
sample dimensionality	15-dim
storage	208 GB
computing time (256 CPUs)	7-8 weeks

Solution approach

- 1 Parallelization
- 2 Model reduction:
 - proper generalized decomposition (PGD)¹
 - sparse grids²

¹Chinesta, Ammar, Leygue and Keunings 2011, ²Delaunay, Lozinski and Owens 2007

Sparse grid combination technique¹

- Approximation of full grid solution $\mathbf{u}_I \in V_I$ as a **combination of coarse full grid solution spaces** $\mathbf{u}_m \in V_m$.
- Multi-indices $\mathbf{m}, \mathbf{l} \in \mathbb{N}^d$ denote discretization accuracy.




FIGURE: Combination technique (left), sparse grid (center) and full grid (right).

¹Griebel, Schneider, Zenger 1992, ²R. and Griebel, submitted to Appl Math & Comput

Sparse grid combination technique¹

- Approximation of full grid solution $\mathbf{u}_I \in V_I$ as a **combination of coarse full grid solution spaces** $\mathbf{u}_m \in V_m$.
- Multi-indices $\mathbf{m}, \mathbf{l} \in \mathbb{N}^d$ denote discretization accuracy.
- Combination technique is **intrinsically parallel**.
- Existing multiscale solver can be **reused**.
- **Numerical result:** Computational effort **reduced by one order of magnitude** in shear and extensional flows².




FIGURE: Combination technique (left), sparse grid (center) and full grid (right).

¹Griebel, Schneider, Zenger 1992, ²R. and Griebel, submitted to Appl Math & Comput

Summary

- ① Simulation of 3D contraction flow with multiscale model:
 - Results compared with experimental measurements from the literature.
 - Experimental phenomena could be reproduced.

Summary

- ① Simulation of 3D contraction flow with multiscale model:
 - Results compared with experimental measurements from the literature.
 - Experimental phenomena could be reproduced.
- ② Reduction of enormous computing time with (twofold parallelism)
 - parallelization (*Atacama* 1248 CPUs).
 - sparse grid combination technique.

Summary

- ① Simulation of 3D contraction flow with multiscale model:
 - Results compared with experimental measurements from the literature.
 - Experimental phenomena could be reproduced.
- ② Reduction of enormous computing time with (twofold parallelism)
 - parallelization (*Atacama* 1248 CPUs).
 - sparse grid combination technique.

Thank you for your attention!

REFERENCES (1 / 3)

- ① C. Le Bris, T. Lelièvre.
Multiscale modelling of complex fluids: a mathematical initiation
Multiscale Modeling and Simulation in Science, 49–137, 2009
- ② M. Laso, H. Öttinger.
Calculation of viscoelastic flow using molecular models
J Non-Newton Fluid Mech, 47:1–20, 1993
- ③ A. Lozinski, C. Chauvière.
A fast solver for Fokker–Planck equation applied to viscoelastic flows calculations
J Comp Phys, 189:607–625, 2003
- ④ H. Warner.
Kinetic theory and rheology of dilute suspensions of finitely extendible dumbbells
Ind Eng Chem Fundam, 11:379–387, 1972
- ⑤ A. Cohen.
A Padé approximant to the inverse Langevin function
Rheologica Acta, 30:270–273, 1991
- ⑥ A. Chorin
Numerical solution of the Navier-Stokes equations
Math. of Comput., 22:745–762, 1968
- ⑦ M. Hulsen, A. Van Heel, B. Van Den Brule
Simulation of viscoelastic flows using Brownian configuration fields
J Non-Newton Fluid Mech, 70:79–101, 1997

REFERENCES (2/3)

- 8 J. von Neumann
Various techniques used in connection with random digits. Monte Carlo methods
Nat Bur Stand, 12:36–38, 1951
- 9 M. Griebel, T. Dornseifer, T. Neunhoeffler
Numerical Simulation in Fluid Dynamics, a Practical Introduction
SIAM, Philadelphia, 1998
- 10 R. Croce, M. Griebel, M.A. Schweitzer
Numerical simulation of bubble and droplet-deformation by a level set approach with surface tension in three dimensions
Int J Numer Methods Fluids, 62:963–993, 2009
- 11 P. Sousa, P. Celho, M. Oliveira, M. Alves
Effect of the contraction ratio upon viscoelastic fluid flow in three-dimensional square-square contractions
Chem Eng Sci, 66:998–1009, 2011
- 12 R. Keunings **On the high Weissenberg number problem**
J Non-Newton Fluid Mech, 20:209–226, 1986
- 13 C. Mangoubi, M. Hulsen, R. Kupferman
Numerical stability of the method of Brownian configuration fields
J Non-Newton Fluid Mech, 157:188–196, 2009
- 14 M. Griebel, A. Rüttgers
Multiscale simulations of 3D viscoelastic flows in a square-square contraction
J Non-Newton Fluid Mech, 205:41–63, 2014

REFERENCES (3/3)

- 15 F. Chinesta, A. Ammar, A. Leygue, R. Keunings
An overview of the proper generalized decomposition with applications in computational rheology
J Non-Newton Fluid Mech, 166:578–592, 2011
- 16 P. Delaunay, A. Lozinski, R. Owens
Sparse tensor-product Fokker-Planck-based methods for nonlinear bead-spring chain models of dilute polymer solutions
CRM Proc, 41:73–89, 2007
- 17 M. Griebel, M. Schneider, C. Zenger
A combination technique for the solution of sparse grid problems
Iterative Meth Lin Alg, 263–281, 1992
- 18 A. Rüttgers, M. Griebel
Multiscale simulation of polymeric fluids using the sparse grid combination technique
submitted to *Appl Math & Comput*